



International Conference

# FROM SEED TO PASTA III

A SUSTAINABLE DURUM WHEAT CHAIN  
FOR FOOD SECURITY AND HEALTHY LIVES



Bologna - Italy, 19-21 September 2018

## P.1.07

### GENOME WIDE ASSOCIATION STUDY REVEALS NOVEL QUANTITATIVE LOCI FOR OSMOTIC ADJUSTMENT UNDER DROUGHT STRESS IN DURUM WHEAT

CONDORELLI G.E.<sup>1</sup>, GROLI E.L.<sup>1</sup>, NEWCOMB M.<sup>2</sup>, MACCAFERRI M.<sup>1</sup>, WARD R.<sup>3</sup>, AND TUBEROSA R.<sup>1</sup>

1) *Department of Agricultural and Food Sciences – DISTAL, University of Bologna, Viale Fanin 44, 40127 Bologna (Italy),*

2) *US Arid Land Agricultural Research Center, Cardon Ln, 21881 Maricopa (United States),*

3) *University of Arizona - Maricopa Agricultural Center (MAC), W Smith Enke Rd 37860, 85138 Maricopa (United States)*

Drought is one of the most devastating stressors limiting crop survival and adaptability, thereby affecting yield loss. Osmotic adjustment (OA) is one of the major components of drought resistance in crops thanks to an active accumulation of solutes in response to a water potential reduction, thereby maintaining turgor. Although yield differences in wheat yield have been correlated to OA capacity, the genetic basis of this trait is unknown.

This work reports the first Genome Wide Association Study (GWAS) in durum wheat for OA as well as for relative water content (RWC), osmotic potential ( $\psi_s$ ) and dry biomass (DB) under full and deficit irrigation conditions. In the field trial in 2018 at the Maricopa Agricultural Center (AZ), 248 durum wheat elite lines and cultivars from worldwide (Durum Panel) were evaluated according to a randomized block design with two replicates. Leaf samples were first collected at flowering under well-irrigated conditions and 10 days later under severe water-stressed conditions. To analyze OA, eight fully expanded flag leaves were sampled from each plot, re-hydrated to full turgor for 8 hours and stored in freezer. The extracted leaf sap was collected for measuring  $\psi_s$  using an osmometer. The resulting osmolality ( $\text{mosmol kg}^{-1}$ ) was converted to osmolarity (MPa) to evaluate OA as  $\psi_s$  (control) -  $\psi_s$  (water-stressed).

The genetic characterization of the Durum Panel was carried out with the Illumina iSelect wheat 90K SNP assay. A Mixed Linear Model (MLM) for GWAS was performed in Tassel in order to identify the genetic intervals harboring loci controlling OA, RWC,  $\psi_s$  and DB. Two major QTL hotspots on chromosomes 6AS and 6AL were found to concurrently affect OA, RWC and DB with a positive effect of OA on both RWC and DB. These results support OA as a prime drought-stress adaptive trait in support of plant production under water-stressed conditions in durum wheat.

ABSTRACT